

Rabies

Annual Epidemiological Report for 2019

Key facts

- For 2019, EU/EEA countries reported five human *Lyssavirus* infections. Four human cases of travel-related rabies were reported by Italy, Latvia, Spain and Norway with exposure in Tanzania, India, Morocco and the Philippines, respectively. One locally-acquired fatal case of *European bat lyssavirus* (EBLV-1) infection was reported by France.

Methods

This report is based on data for 2019 retrieved from The European Surveillance System (TESSy) on 29 October 2020. TESSy is a system for the collection, analysis and dissemination of data on communicable diseases.

For a detailed description of methods used to produce this report, refer to the *Methods* chapter [1].

An overview of the national surveillance systems is available online [2].

A subset of the data used for this report is available through ECDC's online *Surveillance atlas of infectious diseases* [3].

For 2019, 30 EU/EEA countries reported case-based data (Liechtenstein did not report). Twenty-six countries used the EU case definition, three countries used an alternative case definition (Denmark, Germany and Italy) and one country did not specify the case definition used (France). Reporting is compulsory in 29 countries and cases are reported as 'other' in the United Kingdom. Surveillance is comprehensive in all reporting countries and mostly passive.

Epidemiology

For 2019, EU/EEA countries reported five human *Lyssavirus* infections. Four human cases of travel-related rabies were reported by Italy, Latvia, Spain and Norway with exposure in Tanzania, India, Morocco and the Philippines, respectively. The cases reported by Italy, Latvia and Norway were most probably infected through exposure to dogs, while the case reported by Spain was most probably infected through exposure to a cat. One locally-acquired case of *European bat lyssavirus 1* (EBLV-1) infection was reported by France.

For 2014, three cases of rabies were reported by Spain, France and the Netherlands following exposure in Morocco, Mali and India, respectively. For 2015, none were reported. For 2016 and 2017, France reported one travel-related case each year, with exposure in Pakistan and Sri Lanka, respectively. For 2018, one case of travel-related rabies was reported by the United Kingdom. The case was bitten by a cat in Morocco [4,5].

Discussion

In Europe, human rabies is a very rare, vaccine-preventable zoonosis. Once the clinical symptoms have appeared, it is fatal. Very few cases of rabies in humans are reported annually in the EU/EEA and most Member States have not had locally-acquired cases for decades. However, 2019 saw the highest number of rabies cases reported for one year since 2007.

Since 2006, only France and Romania have had reported locally-acquired rabies cases. The case in France resided in the overseas department of French Guiana where he became infected in 2008. In Romania, there was on average one case per year until 2012, the year of the last locally-acquired infection in the country.

Lyssaviruses other than the *rabies lyssavirus* (RABV) have been detected in the EU/EEA and EU neighbouring countries. EBLV-1 and EBLV-2 have bats as their main host and, on rare occasions, they have been linked to infections in other animals and humans [6]. In 2018, 45 EBLV-1 and EBLV-2 bat cases were detected in France, Germany, Hungary, the Netherlands, Poland, Spain and the United Kingdom (UK) [7]. EBLV-1 has been detected in sheep in Denmark [8], in a stone marten in Germany [9], as well as in cats in France [10]. Four human deaths due to EBLV-1 and EBLV-2 have been reported so far in Europe: Ukraine (1977), Russia (1985), Finland (1985) and the UK (2002) [11]. The locally-acquired fatal case of EBLV-1 infection reported by France for 2019 is therefore an unusual, but not unexpected event.

In Europe, bites resulting in rabies exposure are typically from foxes and stray dogs, but also occasionally from raccoons. In many places throughout Asia and Africa, stray dogs are the main source of infections for humans. Illegal importation of pet animals poses a risk of rabies being imported, as reported in France in 2015 [12]. Another source of infection may be through organ transplantation [13]. The re-emergence of animal rabies in northern Italy during the period 2008–2011 and Greece in 2012–2013 underlines the importance of maintaining high levels of awareness [14].

Rabies surveillance data on animals in Europe are available from the ECDC/EFSA summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks [7] and the WHO Collaborating Centre for Rabies Surveillance and Research [15].

Public health implications

It is important to inform the public, especially those travelling to endemic areas, of the risk of contracting rabies if bitten by certain types of mammalian animal hosts, including unvaccinated dogs, foxes, bats and cats [16,17]. Preventive measures include vaccination of domestic carnivores and oral vaccination of wildlife - i.e. red foxes (*Vulpes vulpes*) in Europe.

In the event of exposure to a potentially infected animal, timely prophylaxis is of the utmost importance and knowledge of the epidemiological situation is vital to decide on appropriate post-exposure measures. Treatment consists of wound care, vaccination and passive immunisation with immunoglobulin, if indicated. To be effective, treatment has to be administered as soon as possible after exposure. Specific safety measures should be followed for organ transplantation [15]. People at increased risk of rabies virus infection should consider pre-exposure vaccination.

References

1. European Centre for Disease Prevention and Control. Introduction to the Annual Epidemiological Report Stockholm: ECDC; 2020. Available from: <https://ecdc.europa.eu/en/annual-epidemiological-reports/methods>
2. European Centre for Disease Prevention and Control. Surveillance systems overview [internet, downloadable spreadsheet]. Stockholm: ECDC; 2020. Available from: <https://ecdc.europa.eu/en/publications-data/surveillance-systems-overview-2019>
3. European Centre for Disease Prevention and Control. Surveillance atlas of infectious diseases. Stockholm: ECDC; 2020 [18 Nov 2020]. Available from: <http://atlas.ecdc.europa.eu/public/index.aspx?Dataset=27&HealthTopic=43>
4. Public Health England. Rabies: epidemiology, transmission and prevention. London: PHE; 2018 [updated 2 July 2019]. Available from: <https://www.gov.uk/guidance/rabies-epidemiology-transmission-and-prevention>
5. Public Health England. Public Health England warns travellers of rabies risk. London: PHE; 2018 [updated 12 November 2018]. Available from: <https://www.gov.uk/government/news/public-health-england-warns-travellers-of-rabies-risk>
6. Gossner CM, Mailles A, Aznar I, Dimina E, Echevarria JE, Feruglio SL, et al. Prevention of human rabies: a challenge for the European Union and the European Economic Area. Euro surveillance: bulletin European sur les maladies transmissibles = European communicable disease bulletin. 2020 Sep;25(38).
7. EFSA (European Food Safety Authority) and ECDC (European Centre for Disease Prevention and Control). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2018. EFSA Journal. 2019. 18 Nov 2020;17(12):5926.
8. Tjornehoj K, Fooks AR, Agerholm JS, Ronsholt L. Natural and experimental infection of sheep with European bat lyssavirus type-1 of Danish bat origin. J Comp Pathol. 2006 Feb-Apr;134(2-3):190-201.
9. Muller T, Cox J, Peter W, Schafer R, Johnson N, McElhinney LM, et al. Spill-over of European bat lyssavirus type 1 into a stone marten (Martes foina) in Germany. J Vet Med B Infect Dis Vet Public Health. 2004 Mar;51(2):49-54.
10. Dacheux L, Larrous F, Mailles A, Boisseleau D, Delmas O, Biron C, et al. European bat *Lyssavirus* transmission among cats, Europe. Emerging Infectious Diseases. 2009 Feb;15(2):280-4.
11. Fooks AR, Brookes SM, Johnson N, McElhinney LM, Hutson AM. European bat lyssaviruses: an emerging zoonosis. Epidemiology and Infection. 2003 Dec;131(3):1029-39.
12. Rabies confirmed in an illegally imported dog in France. The Veterinary Record. 2015 May 30;176(22):558.
13. Maier T, Schwarting A, Mauer D, Ross RS, Martens A, Kliem V, et al. Management and outcomes after multiple corneal and solid organ transplantations from a donor infected with rabies virus. Clinical Infectious Diseases: an official publication of the Infectious Diseases Society of America. 2010 Apr 15;50(8):1112-9.
14. Tsiodras S, Dougas G, Baka A, Billinis C, Doudounakis S, Balaska A, et al. Re-emergence of animal rabies in northern Greece and subsequent human exposure, October 2012 - March 2013. Euro surveillance: bulletin European sur les maladies transmissibles = European communicable disease bulletin. 2013 May 2;18(18):20474.
15. World Health Organization. Rabies - Prevention Geneva: WHO; 2020 [cited 18 Nov 2020]. Available from: http://www.who.int/rabies/about/home_prevention/en/
16. Cliquet F, Picard-Meyer E, Robardet E. Rabies in Europe: what are the risks? Expert Review of Anti-infective Therapy. 2014 Aug;12(8):905-8.
17. Public Health England. Rabies risks in terrestrial animals by country. London: PHE; 2019 [updated 1 May 2019; cited 18 October 2019]. Available from: <https://www.gov.uk/government/publications/rabies-risks-by-country/rabies-risks-in-terrestrial-animals-by-country>