# Tuberculosis surveillance and monitoring in Europe 2019





2017 data

# **Germany**

Total population at 23 April 2018 by EUROSTAT: 82 521 65

#### TB case notifications, 2017

Total number of cases	5 486		
Notification rate per 100 000	6.6		
New <sup>a</sup> and relapses	5 353		
New <sup>a</sup> and relapses notification rate per 100 000	6.5		
Pulmonary	3 892	(70.9%)	
of which microscopy-positive	1804	(46.4%)	
of which laboratory-confirmed	3 090	(79.4%)	
Laboratory-confirmed TB cases	3 978	(72.5%)	
Mean age of new native TB cases	65.4	65.4 years	
Mean age of new foreign TB cases	40.8	40.8 years	
Foreign origin of all TB cases	3 820	(69.6%)	
New (not previously treated)	5 252	(95.7%)	

<sup>&</sup>lt;sup>a</sup> Cases with unknown previous treatment included in new cases.

## Drug resistance surveillance and TB/HIV coinfection, 2017

Completeness of DRS data <sup>a</sup>	Yes	
Completeness of HIV datab	-	
Case-linked data reporting	Yes	
Cases with DST results	3 555	(89.4%)
Estimated RR-TB among notified pulmonary cases N, (best-low-high)	140-65-210	
Pulmonary MDR-TB cases notified	87	(3.1%)
of which XDR-TB cases	4	(5.7%)
Notified MDR-TB	105	(3.0%)
of which XDR-TB cases	4	(4.9%)
TB cases tested for HIV	-	-
HIV-positive TB cases	_	_
of these on ART	-	-


 $<sup>^</sup>a$  National coverage 100% or culturing  $\ge$  90%, C+/all TB cases > 50%, DST done for C+ > 75%, EQA  $\ge$  95%.  $^b$  More than 50% of TB cases with reported HIV status.

Treatment outcome monitoring


Geographical coverage	Na	tional		
Outcome cohort	New culture positive pulmonary TB cases notified in 2016 <sup>a</sup>		All MDR-TB cases notified in 2015 <sup>b</sup>	
Case-linked data reporting	Yes			
Cases notified	199		110	
Success	118	(59.3%)	44	(40.0%)
Died	17	(8.5%)	5	(4.5%)
Failed	0	(0.0%)	0	(0.0%)
Lost to follow-up	2	(1.0%)	7	(6.4%)
Still on treatment	12	(6.0%)	11	(10.0%)
Not evaluated	50	(25.1%)	43	(39.1%)

 $<sup>^{\</sup>rm a}$  Treatment outcome as presented is treatment outcome after 12 months.  $^{\rm b}$  Treatment outcome as presented is treatment outcome after 24 months.


TB notification rates by treatment history, 2008-2017



## New and relapsed TB cases – notification rates by age group, 2008–2017




### TB cases by geographical origin, 2008-2017




### TB/HIV coinfection, 2008-2017

Data not available

## MDR-TB cases by previous treatment history, 2008-2017



## Treatment outcome, new culture-confirmed pulmonary TB cases, 2007–2016



Source: European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Tuberculosis surveillance and monitoring in Europe 2019 – 2017 data. Stockholm: ECDC; 2019