

Summary

Week 14/2023 (03 April–09 April 2023)

- The percentage of all sentinel primary care specimens from patients presenting with ILI or ARI symptoms that tested positive for an influenza virus fluctuated to 15% in week 14/2023 from 16% in the previous week and remaining above the epidemic threshold (10%).
- 11 of 42 countries or areas reported medium intensity and 16 of 41 countries across the Region reported widespread activity.
- Of the 29 countries that reported sentinel primary care specimen influenza virus positivity above the 10% epidemic threshold, only Estonia and Slovakia reported activity above 40%.
- Influenza type A and type B viruses were detected in sentinel and non-sentinel surveillance, with influenza type B predominating in both systems.
- Hospitalized patients with confirmed influenza virus infection were reported from ICU (with higher proportions of type B viruses) and SARI surveillance (with higher proportions of type B viruses). Four countries or areas reported influenza virus positivity rates above 10% in SARI surveillance (Lithuania, Albania, Serbia and Ukraine).

2022–2023 season overview

- The seasonal epidemic activity threshold of 10% positivity in sentinel specimens was first crossed in week 45/2022.
- Following a peak at week 51/2022 with 39% positivity, influenza activity had been decreasing across the Region until week 4/2023 when it reached 22% positivity before rising again to fluctuate around 25% positivity between weeks 6 and 11/2023 before decreasing again to 15% positivity for week 14/2023.
- Overall this season, influenza A(H3) viruses have dominated in sentinel primary care specimens, however higher circulation of A(H1)pdm09 and type B viruses was observed starting from week 50/2022 and week 2/2023, respectively. A similar prevalence of A(H1)pdm09 and A(H3) viruses was detected in non-sentinel specimens.
- Both influenza type A and type B viruses have been detected in hospitalized patients in ICU and other wards and influenza A(H1)pdm09 viruses have dominated among SARI patients.
- Virus type and subtype prevalence by country and surveillance system has been variable across the season.
- Few B/Yamagata viruses reported by different countries across the season have been further investigated and were proven to be LAIV infections.

Other news

- RSV is another respiratory virus that causes acute respiratory disease, mainly among young infants and the elderly, often mild but frequently severe among children less than 1 year of age and frail elderly. High levels of RSV have been circulating across the Region since week 40/2022, with overall positivity amongst patients in primary care with acute respiratory illness decreasing after a peak at 18% positivity in week 47/2022 to 4% for week 14/2023. More information on the risk of RSV infections can be found [here](#).

For more information about the SARS-CoV-2 situation in the WHO European Region visit:

- WHO website: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
- ECDC website: <https://www.ecdc.europa.eu/en/novel-coronavirus-china>

Qualitative indicators

For week 14/2023, of 42 countries and areas reporting on intensity of influenza activity, 12 reported baseline-intensity (across the Region), 19 reported low-intensity (across the Region) and 11 reported medium-intensity (across the Region) (Fig. 1).

Of 41 countries and areas reporting on geographic spread of influenza viruses, 3 reported no activity (Georgia, Kazakhstan and Kyrgyzstan), 9 reported sporadic spread (eastern, northern and southern Region), 5 reported local spread (France, Malta, Serbia, Slovakia and Kosovo (in accordance with UN Security Council Resolution 1244 (1999))), 8 reported regional spread (Albania, Austria, Bosnia and Herzegovina, Czechia, Lithuania, Republic of Moldova, Romania and Ukraine) and 16 reported widespread activity (across the Region) (Fig. 2).

Figure 1. Intensity of influenza activity in the European Region, week 14/2023





Figure 2. Geographic spread of influenza viruses in the European Region, week 14/2023



For interactive maps of influenza intensity and geographic spread, see the [Flu News Europe website](#).

Please note:


- Assessment of the intensity of activity indicator includes consideration of ILI or ARI rates. These ILI or ARI rates might be driven by respiratory infections other than influenza virus, including SARS-CoV-2, leading to observed increases in the absence of influenza virus detections.
- Assessment of intensity and geographic spread indicators includes consideration of sentinel and non-sentinel influenza virus detection data. Non-sentinel influenza virus detections, often higher, might translate into reporting of elevated geographic spread even in the absence of sentinel detections and/or low intensity of activity measured by ILI and ARI incidence.

Influenza positivity

For the European Region, influenza virus positivity in sentinel primary care specimens fluctuated from 16% in the previous week to 15% in week 14/2023. Seasonal activity started in week 45/2022 when positivity crossed above the epidemic threshold set at 10%. The current seasonal influenza epidemic started earlier than in the four previous seasons, ranging from week 47 (2019/20 season) to 49 (2021/22 season). Positivity reached a peak in week 51/2022 at 39% which was earlier than in the four previous seasons, ranging from week 52 (2021/22 season) to 5 (2018/19 and 2019/20). Across the Region, influenza activity decreased to 22% up to week 4/2023, then fluctuated around 25% between weeks 6 and 11/2023 and decreased again to 15% in week 14/2023 (Fig. 3).

Figure 3. Influenza virus positivity in sentinel-source specimens by week, WHO European Region, seasons 2021/2022, 2019/2020, 2018/2019, 2017/2018 and 2022/2023

Influenza positivity in sentinel-source specimens by week - WHO Europe



External data sources

Mortality monitoring:

The EuroMOMO report can be found here: <https://www.euromomo.eu/>

Please refer to the EuroMOMO website for a cautionary note relating to interpretation of these data.

Primary care data

Syndromic surveillance data

Of the countries and areas in which thresholds for ILI activity are defined, countries in eastern (n=6; Azerbaijan, Georgia, Kazakhstan, Kyrgyzstan, Russian Federation and Uzbekistan), northern (n=3; Denmark, Estonia and Lithuania), southern (n=3; Croatia, Greece and Serbia) and western (n=3; Hungary, Luxembourg and Poland) areas of the European Region reported activity above baseline levels.

Of the countries and areas in which thresholds for ARI activity are defined, countries in eastern (n=2; Kazakhstan and Uzbekistan) and northern (n=1; Latvia) areas of the European Region reported activity above baseline levels.

Please note:

- Assessment of the syndromic surveillance data of ILI or ARI rates might be driven by respiratory infections other than influenza virus, including SARS-CoV-2, leading to observed increases in the absence of influenza virus detections. The thresholds mentioned are related to the Moving Epidemic Method (MEM) method and based on historic ILI/ARI data.

Viruses detected in sentinel-source specimens (ILI and ARI)

For week 14/2023, 328 (15%) of 2 118 sentinel specimens tested positive for an influenza virus; 83% were type B and 17% were type A. Of 14 subtyped A viruses, 86% were A(H1)pdm09 and 14% A(H3). All 53 type B viruses ascribed to a lineage were B/Victoria (Fig. 4 and Table 1). Of 29 countries and areas across the Region that each tested at least 10 sentinel specimens in week 14/2023, 15 reported a rate of influenza virus detections above 10% (median 25%; range 12% - 46%): Estonia (46%), Slovakia (45%), Hungary (38%), France (29%), Poland (29%), Germany (27%), Romania (25%), Slovenia (25%), Luxembourg (24%), Kosovo (in accordance with Security Council resolution 1244 (1999)) (24%), Norway (23%), Armenia (19%), Austria (15%), Ukraine (14%) and Italy (12%).

For the season to date, 27 003 (23%) of 116 620 sentinel specimens tested positive for an influenza virus. More influenza type A (n=19 291, 71%) than type B (n=7 712, 29%)

viruses have been detected. Of 15 613 subtyped A viruses, 10 023 (64%) were A(H3) and 5 590 (36%) were A(H1)pdm09. All 2 272 influenza type B viruses ascribed to a lineage were B/Victoria (71% of type B viruses were reported without a lineage) (Fig. 4 and Table 1).

Details of the distribution of viruses detected in non-sentinel-source specimens are presented in the **virus characteristics** section.

Figure 4. Influenza virus positivity and detections by type, subtype/lineage – sentinel sources, WHO European Region, season 2022/2023

Influenza virus positivity and detections by type, subtype/lineage and week - WHO Europe, season 2022/2023




Table 1. Influenza virus detections in sentinel source specimens by type and subtype for week 14/2023 and cumulatively for the season

Sentinel	Current Week (14)		Season 2022-2023	
Virus type and subtype	Number	% ^a	Number	% ^a
Influenza A	55	16.8	19 291	71.4
A(H1)pdm09	12	85.7	5 590	35.8
A(H3)	2	14.3	10 023	64.2
A not subtyped	41	-	3 678	-
Influenza B	273	83.2	7 712	28.6
B/Victoria lineage	53	100	2 272	100
B/Yamagata lineage	0	0	0	0
Unknown lineage	220	-	5 440	-
Total detections (total tested)	328 (2 118)	15.4	27 003 (116 620)	23.2

^a For influenza type percentage calculations, the denominator is total detections; for subtype and lineage, it is total influenza A subtyped and total influenza B lineage determined, respectively; for total detections, it is total tested.

External data sources

Influenzanet collects weekly data on symptoms in the general community from different participating countries across the EU/EEA. Please refer to the website for information for this week.

Hospital surveillance

A subset of Member States and areas monitors severe disease related to influenza virus infection by surveillance of 1) hospitalized laboratory-confirmed influenza cases in ICUs, or other wards, or 2) severe acute respiratory infections (SARI).

Laboratory-confirmed hospitalized cases

1.1) Hospitalized laboratory-confirmed influenza cases - Intensive care units (ICUs)

For week 14/2023, 7 laboratory-confirmed influenza cases were reported from ICU wards (in Czechia (n=1), France (n=1) and Sweden (n=5)). Both influenza type A viruses (n=2) and type B viruses (n=5) were detected. No viruses were ascribed to a subtype or lineage (Fig. 5 and 6).

Since week 40/2022, 2 695 influenza type A (90%) and 289 type B (10%) viruses were detected (in Czechia (n=141), France (n=930), Ireland (n=151), Sweden (n=258) and United Kingdom (England) (n=1 504)). Of 487 subtyped influenza A viruses, 53% were A(H3) and 47% were A(H1)pdm09. No influenza B viruses were ascribed to a lineage. Of 1 470 cases with known age, 695 were 15-64 years old, 593 were 65 years and older, 114 were 0-4 years old and 68 were 5-14 years old.

Figure 5. Number of laboratory-confirmed hospitalized influenza cases in intensive care units (ICU) by week of reporting, WHO European Region, season 2022/2023





Figure 6. Distribution of influenza virus types, subtypes/lineages by age group in intensive care units (ICU), WHO European Region, season 2022/2023

Distribution of virus types, subtypes/lineages by age group in intensive care units (ICU) - WHO Europe, season 2022/2023



1.2) Hospitalized laboratory-confirmed influenza cases – other wards

For week 14/2023, there were no reports of hospitalized laboratory-confirmed influenza cases in other wards (Fig. 7 and 8). No viruses were ascribed to a subtype or lineage (Fig. 7 and 8).

Since week 40/2022, 3807 influenza type A viruses and 179 influenza type B viruses were detected from Czechia (n=169) and Ireland (n=3 817). Of 397 subtyped influenza A viruses, 63% (n=251) were A(H1)pdm09 and 37% (n=146) A(H3). No influenza B viruses were ascribed to a lineage. The 3 986 cases with known age fell in 4 age groups: 1712 were 65 years and older, 1374 were 15-64 years old, 499 were 0-4 years old and 401 were 5-14 years old.

Figure 7. Number of laboratory-confirmed hospitalized influenza cases in wards other than intensive care units (non-ICU) by week of reporting, WHO European Region, season 2022/2023





Figure 8. Distribution of influenza virus types, subtypes/lineages by age group in wards other than intensive care units (non-ICU), WHO European Region, season 2022/2023



Severe acute respiratory infection (SARI)-based hospital surveillance

For week 14/2023, 2 709 SARI cases were reported by 19 countries or areas: Albania, Belarus, Bosnia and Herzegovina, Georgia, Germany, Ireland, Kazakhstan, Kyrgyzstan, Lithuania, Malta, North Macedonia, Republic of Moldova, Romania, Russian Federation, Serbia, Spain, Ukraine, Uzbekistan and Kosovo (in accordance with Security Council resolution 1244 (1999)). Of 987 specimens tested for influenza viruses, 6% (n=59) were positive (Fig. 9). Of these, influenza type B viruses (n=49, 83%) were detected more

frequently than influenza type A viruses (n=10, 17%). Of 3 subtyped influenza type A viruses, 2 were A(H1)pdm09 and 1 was A(H3). All 2 type B viruses ascribed to a lineage were B/Victoria. Of 14 countries and areas across the Region that each tested at least 10 specimens, 4 reported positivity rates above 10%: Lithuania (40%), Albania (20%), Serbia (17%) and Ukraine (15%).

For the season, 129 697 SARI cases were reported by 26 countries or areas (Albania, Armenia, Belarus, Belgium, Bosnia and Herzegovina, Croatia, Georgia, Germany, Ireland, Kazakhstan, Kyrgyzstan, Lithuania, Malta, Montenegro, North Macedonia, Republic of Moldova, Romania, Russian Federation, Serbia, Slovakia, Spain, Türkiye, Turkmenistan, Ukraine, Uzbekistan and Kosovo (in accordance with Security Council resolution 1244 (1999))). For SARI cases testing positive for influenza virus since week 40/2022, type A viruses have been the most common (n=3 448, 69%) and of these 2 766 were subtyped: 2 050 (74%) were infected by A(H1)pdm09 viruses and 716 (26%) were infected by A(H3) viruses. Of those influenza B viruses that have been ascribed to a lineage (n=391, 31%), all were B/Victoria (Fig. 10).

Figure 9. Number of severe acute respiratory infection (SARI) cases (bar) and positivity for influenza virus and SARS-CoV-2 (line) by week, WHO European Region, season 2022/2023





Figure 10. Influenza virus detections by type, subtype/lineage from severe acute respiratory infection (SARI), WHO European Region, season 2022/2023



Virus characteristics

Details of the distribution of viruses detected in sentinel-source specimens can be found in the **Primary care data** section.

Non-sentinel virologic data

For week 14/2023, 2 429 of 44 097 specimens from non-sentinel sources (such as hospitals, schools, primary care facilities not involved in sentinel surveillance, or nursing homes and other institutions) tested positive for an influenza virus; 754 (31%) were type A and 1 675 (69%) were type B. Of 67 subtyped A viruses, 50 (75%) were A(H1)pdm09 and 17 (25%) A(H3). Of 83 type B viruses ascribed to a lineage, 74 (89.2%) were B/Victoria and 9 (10.8%) were B/Yamagata reported by a single country. The B/Yamagata reports are under investigation (Fig. 11 and Table 2).

For the season to date, more influenza type A (n=191 407, 76%) than type B (n=59 143, 24%) viruses have been detected. Of 55 749 subtyped A viruses, 30 774 (55%) were A(H1)pdm09 and 24 975 (45%) were A(H3). Of 4 379 influenza type B viruses ascribed to a lineage, 4 370 were B/Victoria and 9 were B/Yamagata (93% of type B viruses were reported without a lineage). These reported B/Yamagata viruses are under investigation. The confirmed B/Yamagata LAIV infections are not included in the season's count (Fig. 11 and Table 2).

Figure 11. Influenza detections by type, subtype/lineage and week, non-sentinel sources, WHO European Region, season 2022/2023

Influenza virus detections by type, subtype/lineage and week - WHO Europe, season 2022/2023




Table 2. Influenza virus detections in non-sentinel-source specimens by type and subtype, week 14/2023 and cumulatively for the season

* Due to a reporting error, this figure cannot be shown at this time.

^a For type percentage calculations, the denominator is total detections; for subtype and lineage, it is total influenza A subtyped and total influenza B lineage determined, respectively; as not all countries have a true non-sentinel testing denominator, no percentage calculations for total tested are shown.

Genetic characterization

Of the 2 345 genetically characterized A(H1)pdm09 viruses up to week 14/2023, 1 329 were attributed to clade 6B.1A.5a.2, of which 665 (50%) were represented by A/Norway/25089/2022, 631 (47%) by A/Sydney/5/2021 and 33 (2%) by A/Victoria/2570/2019. 5 (<1%) were attributed to clade 6B.1A.5a.1 represented by A/Guangdong-Maonan/SWL1536/2019. 1 011 (43%) viruses could not be attributed to a pre-defined subgroup in the guidance.

Among the 2 435 A(H3) viruses characterized up to week 14/2023, 2 315 were attributed to clade 3C.2a1b.2a.2, of which 1 423 (61%) were represented by A/Bangladesh/4005/2020, 746 (32%) by A/Slovenia/8720/2022, 146 (6%) by A/Darwin/9/2021. Only 3 viruses were ascribed to clade 3C.2a1b.1a represented by A/Denmark/3264/2019. 117 (5%) viruses could not be attributed to a pre-defined subgroup in the guidance.

Up to week 14/2023, 870 B/Victoria viruses were characterized, 554 (64%) of which were attributed to clade V1A.3a.2 represented by B/Austria/1359417/2021. 316 (36%) viruses could not be attributed to a pre-defined subgroup in the guidance.

Table 3. Number of influenza viruses attributed to genetic groups, cumulative for the season, WHO European Region

Number of influenza viruses attributed to genetic groups, cumulative for the season - WHO Europe

<div> <div>Virus Type</div> <div>Virus Subtype</div> <div>Genetic charact...</div> </div>		Number of influenza viruses attributed to genetic groups
		2022/2023
Total		5 650
Influenza A		4 780
A(H1)pdm09		2 345
A(H1)pdm09_SubgroupNotListed *		1 011
A/Guangdong-Maonan/SWL1536/2019(H1N1)pdm09_6B.1A.5a.1		5
A/Norway/25089/2022(H1N1)pdm09_6B.1A.5a.2		665
A/Sydney/5/2021(H1N1)pdm09_6B.1A.5a.2		631
A/Victoria/2570/2019(H1N1)pdm09_6B.1A.5a.2		33
A(H3)		2 435
A(H3)_SubgroupNotListed *		117
A/Bangladesh/4005/2020(H3)_3C.2a1b.2a.2		1 423
A/Darwin/9/2021(H3)_3C.2a1b.2a.2		146
A/Denmark/3264/2019(H3N2)_3C.2a1b.1a		3
A/Slovenia/8720/2022(H3)_3C.2a1b.2a.2		746
Influenza B		870
B/Vic		870
B/Austria/1359417/2021(Victoria lineage_1A.3a.2)		554
BVic_SubgroupNotListed *		316

* No Clade: not attributed to a pre-defined clade and SubgroupNotListed: attributed to recognised group in current guidance but not listed here

© World Health Organization 2022.

© European Centre for Disease Prevention and Control 2022.

Reproduction is authorised, provided the source is acknowledged.



Currently, [WHO Europe and ECDC's February](#) virus characterization report is available and describes available data from circulating viruses for the early weeks of the 2022-2023 influenza season.

Antiviral susceptibility testing

Up to week 14/2023, 4 149 viruses were assessed for susceptibility to neuraminidase inhibitors (1 416 A(H1)pdm09, 1 305 A(H3) and 714 B viruses only genotypically and 321 A(H3), 250 A(H1)pdm09 and 143 B viruses phenotypically and some of these also genotypically), and 3 035 viruses were assessed for susceptibility to baloxavir marboxil (1 611 A(H3), 827 A(H1)pdm09 and 597 B viruses genotypically). Phenotypically and/or genotypically, 5 A(H1)pdm09 viruses showing (highly) reduced inhibition by oseltamivir and normal inhibition by zanamivir were identified of which 4 were reported to carry reduced inhibition markers, 3 with NA-H275Y (1 confirmed phenotypically) and 1 with NA-D199G (confirmed phenotypically), and for 1 the amino acid change was not reported. Phenotypically, one A(H3) virus showing highly reduced inhibition by oseltamivir was identified. Genotypically no markers associated with reduced susceptibility for baloxavir marboxil were identified.

Vaccine

Recently published results from a controlled, randomised trial in UK concluded that concomitant vaccination with one of two SARS-CoV-2 vaccines (ChAdOx1 or BNT162b2) plus an age-appropriate influenza vaccine raised no safety concerns and preserves [antibody responses](#) to both vaccines.

[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)02329-1/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)02329-1/fulltext)

Available vaccines in Europe <https://www.ecdc.europa.eu/en/seasonal-influenza/prevention-and-control/vaccines/types-of-seasonal-influenza-vaccine>

Vaccine composition

On 24 February 2023, WHO published recommendations for the components of influenza vaccines for use in the 2023-2024 northern hemisphere influenza season:

The WHO recommends that quadrivalent vaccines for use in the 2023-2024 influenza season in the northern hemisphere contain the following:

Egg-based Vaccines

- an A/Victoria/4897/2022 (H1N1)pdm09-like virus;
- an A/Darwin/9/2021 (H3N2)-like virus;
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus; and
- a B/Phuket/3073/2013 (B/Yamagata lineage)-like virus.

Cell culture- or recombinant-based Vaccines

- an A/Wisconsin/67/2022 (H1N1)pdm09-like virus;
- an A/Darwin/6/2021 (H3N2)-like virus;
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus; and
- a B/Phuket/3073/2013 (B/Yamagata lineage)-like virus.

The WHO recommends that trivalent vaccines for use in the 2023-2024 influenza season in the northern hemisphere contain the following:

Egg-based vaccines

- an A/Victoria/4897/2022 (H1N1)pdm09-like virus;
- an A/Darwin/9/2021 (H3N2)-like virus; and
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus.

Cell culture- or recombinant-based vaccines

- an A/Wisconsin/67/2022 (H1N1)pdm09-like virus;
- an A/Darwin/6/2021 (H3N2)-like virus; and
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus

The full report is published [here](#).

On 23 September 2022, WHO published recommendations for the components of influenza vaccines for use in the 2023 southern hemisphere influenza season:

Egg-based Vaccines

- an A/Sydney/5/2021 (H1N1)pdm09-like virus;
- an A/Darwin/9/2021 (H3N2)-like virus;
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus; and
- a B/Phuket/3073/2013 (B/Yamagata lineage)-like virus.

Cell- or recombinant-based Vaccines

- an A/Sydney/5/2021 (H1N1)pdm09-like virus;
- an A/Darwin/6/2021 (H3N2)-like virus;
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus; and
- a B/Phuket/3073/2013 (B/Yamagata lineage)-like virus.

It is recommended that **trivalent influenza vaccines** for use in the 2022 southern hemisphere influenza season contain the following:

Egg-based vaccines

- an A/Sydney/5/2021 (H1N1)pdm09-like virus;
- an A/Darwin/9/2021 (H3N2)-like virus; and
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus.

Cell- or Recombinant-based vaccines

- an A/Sydney/5/2021 (H1N1)pdm09-like virus;
- an A/Darwin/6/2021 (H3N2)-like virus; and
- a B/Austria/1359417/2021 (B/Victoria lineage)-like virus

The full report is published [here](#).

Acknowledgements

This weekly update was prepared by an editorial team at the European Centre for Disease Prevention and Control (Cornelia Adlhoch, Maja Vukovikj, and Edoardo Colzani) and the WHO Regional Office for Europe (Margaux Meslé, Piers Mook and Richard Pebody). It was reviewed by experts from the network (Adam Meijer, National Institute for Public Health and the Environment (RIVM), the Netherlands); Monica Galiano, WHO Collaborating Centre for Reference and Research on Influenza, Francis Crick Institute, United Kingdom.

Maps and commentary do not represent a statement on the legal or border status of the countries and territories shown.

All data are up to date on the day of publication. Past this date, however, published data should not be used for longitudinal comparisons, as countries retrospectively update their databases. The WHO Regional Office for Europe is responsible for the accuracy of the Russian translation.

Suggested citation: European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Flu News Europe, Joint ECDC–WHO weekly influenza update, week 14/2023.

Tables and figures should be referenced:

European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Flu News Europe, Joint ECDC–WHO weekly influenza update, week 14/2023.

© World Health Organization 2023

© European Centre for Disease Prevention and Control 2023

Reproduction is authorized, provided the source is acknowledged.