Tuberculosis surveillance and monitoring in Europe 2018.





2016 data

Belgium

Total population at 22 September 2017 by EUROSTAT: 11311117

Tuberculosis case notifications, 2016

Total number of cases	1047	
Notification rate per 100 000	9.3	
New* & relapses	986	
New* & relapses notification rate per 100 000	8.7	
Pulmonary	715	(68.3%)
of which microscopy-positive	331	(46.3%)
of which laboratory confirmed	593	(82.9%)
Laboratory-confirmed TB cases	801	(76.5%)
Mean age of new native TB cases	45.2 years	
Mean age of new foreign TB cases	35.5 years	
Foreign origin of all TB cases	547	(52.2%)
New (not previously treated)	757	(72.3%)

^{*} Cases with unknown previous treatment included in new cases.

Drug resistance surveillance & TB-HIV co-infection, 2016

Completeness of DRS data*	١	/es	
Completeness of HIV data**		No	
Case-linked data reporting	}	Yes	
Cases with DST results	769	(96.0%)	
Estimated RR-TB among notified pulmonary cases N, (best-low-high)	16-6-26		
Pulmonary MDR TB cases notified	9	(1.6%)	
of which XDR TB cases	0	(0.0%)	
Notified MDR TB	10	(1.3%)	
of which XDR TB cases	0	(0.0%)	
TB cases tested for HIV	479	(45.7%)	
HIV-positive TB cases	33	(6.9%)	
of these on antiretroviral therapy	-	-	


^{*} National coverage 100% or culturing 90%, C+/All TB

Treatment outcome monitoring


Geographical coverage	Na	tional		
Outcome cohort	New culture positive pulmonary TB cases notified in 2015*		All MDR TB cases notified in 2014**	
Case-linked data reporting	,	/es		
Cases notified	422		11	
Success	351	(79.4%)	6	(54.5%)
Died	34	(7.7%)	1	(9.1%)
Failed	0	(0.0%)	0	(0.0%)
Lost to follow-up	33	(7.5%)	0	(0.0%)
Still on treatment	12	(2.7%)	4	(36.4%)
Not evaluated	12	(2.7%)	0	(0.0%)

^{*} Treatment outcome as presented is treatment outcome after 12 months.


Tuberculosis notification rates by treatment history, 2007–2016




New and relapsed TB cases – notification rates by age group, 2007–2016




Tuberculosis cases by geographical origin, 2007-2016




TB-HIV co-infection, 2007-2016



MDR TB cases by previous treatment history, 2007–2016



Treatment outcome, new culture-confirmed pulmonary TB cases, 2006–2015



Source: European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Tuberculosis surveillance and monitoring in Europe 2018 – 2016 data. Stockholm: ECDC; 2018

^{**} More than 50% of TB cases with reported HIV status

^{**} Treatment outcome as presented is treatment outcome after 24 months.