

References

1. Galati AB, Galvis-Ovallos F, editors. Gaps in phlebotomine distribution and their taxonomic implications. 10th International Symposium on Phlebotomine Sand flies; 2019; San Cristobal, Galapagos, Ecuador: Universidad de San Francisco de Quito.
2. Alten B, Maia C, Afonso MO, Campino L, Jimenez M, Gonzalez E, et al. Seasonal dynamics of phlebotomine sand fly species proven vectors of Mediterranean leishmaniasis caused by *Leishmania infantum*. PLoS Negl Trop Dis. 2016 Feb;10(2).
3. Kasap OE, Votycka J, Alten B. The distribution of the *Phlebotomus major* complex (Diptera: Psychodidae) in Turkey. Acta Trop. 2013 Sep;127(3):204-11.
4. Medlock JM, Hansford KM, Van Bortel W, Zeller H, Alten B. A summary of the evidence for the change in European distribution of phlebotomine sand flies (Diptera: Psychodidae) of public health importance. J Vector Ecol. 2014 Jun;39(1):72-7.
5. Ayhan N, Charrel RN. Of phlebotomines (sand flies) and viruses: a comprehensive perspective on a complex situation. Current Opinion in Insect Science. 2017 Aug;22:117-24.
6. Feliciangeli MD. Natural breeding places of phlebotomine sand flies (vol 18, pg 71, 2004). Med Vet Entomol. 2004 Dec;18(4):453-4.
7. Munoz C, Risueno J, Yilmaz A, Perez-Cutillas P, Goyena E, Ortuno M, et al. Investigations of *Phlebotomus perniciosus* sand flies in rural Spain reveal strongly aggregated and gender-specific spatial distributions and advocate use of light-attraction traps. Med Vet Entomol. 2018 Jun;32(2):186-96.
8. Molina R, Jimenez MI, Cruz I, Iriso A, Martin-Martin I, Sevillano O, et al. The hare (*Lepus granatensis*) as potential sylvatic reservoir of *Leishmania infantum* in Spain. Vet Parasitol. 2012 Nov 23;190(1-2):268-71.
9. Koch LK, Kochmann J, Klimpel S, Cunze S. Modeling the climatic suitability of leishmaniasis vector species in Europe. Sci Rep. 2017 Oct 17;7.
10. Rioux JA, Golvan YJ, Croset H, Tour S, Houin R, Abonnenc E, et al. Epidémiologie des Leishmanioses dans le Sud de la France. Paris: Institut National de la Santé et de la Recherche Médicale; 1969. p. 223.
11. Ballart C, Baron S, Alcover MM, Portus M, Gallego M. Distribution of phlebotomine sand flies (Diptera: Psychodidae) in Andorra: First finding of *P. perniciosus* and wide distribution of *P. ariasi*. Acta Trop. 2012 Apr;122(1):155-9.
12. Merino-Espinosa G, Corpas-Lopez V, Callejon-Fernandez R, Porcel-Rodriguez L, Diaz-Saez V, Gallego M, et al. Differential ecological traits of two *Phlebotomus sergenti* mitochondrial lineages in southwestern Europe and their epidemiological implications. Trop Med Int Health. 2016 May;21(5):630-41.
13. Munoz C, Martinez-de la Puente J, Figuerola J, Perez-Cutillas P, Navarro R, Ortuno M, et al. Molecular xenomonitoring and host identification of *Leishmania* sand fly vectors in a Mediterranean periurban wildlife park. Transbound Emerg Dis. 2019 Nov;66(6):2546-61.
14. Ready PD. Leishmaniasis emergence in Europe. Eurosurveillance. 2010 Mar 11;15(10):29-39.
15. Semenza JC, Suk JE. Vector-borne diseases and climate change: a European perspective. FEMS Microbiol Lett. 2018 Jan;365(2).
16. Fischer D, Thomas SM, Beierkuhnlein C. Modelling climatic suitability and dispersal for disease vectors: the example of a phlebotomine sandfly in Europe. Spatial Statistics 2011: Mapping Global Change. 2011 2011;7:164-9.
17. Fischer D, Moeller P, Thomas SM, Naucke TJ, Beierkuhnlein C. Combining climatic projections and dispersal ability: a method for estimating the responses of sandfly vector species to climate change. PLoS Negl Trop Dis. 2011 Nov;5(11).
18. Trajer AJ, Bede-Fazekas A, Hufnagel L, Horvath L, Bobvos J, Paldy A. The effect of climate change on the potential distribution of the European *Phlebotomus* species. Appl Ecol Environ Res. 2013 2013;11(2):189-208.
19. Lane RP. Medical insects and arachnids. In: Lane R, Crosskey R, editors. Medical insects and arachnids. London: Chapman and Hall; 1993. p. 78-119.
20. Service M. Phlebotomine sand-flies. In: Service M, editor. Medical Entomology for Students 4th Edition. Cambridge: Cambridge University Press; 2008.
21. Kamhawi S. The biological and immunomodulatory properties of sand fly saliva and its role in the establishment of *Leishmania* infections. Microb Infect. 2000 Nov;2(14):1765-73.
22. Lawyer PG, Perkins PV. Leishmaniasis and trypanosomiasis. In: Eldridge B, Edman J, editors. Medical Entomology. Dordrecht: Kluwer Academic Publishers 2000.
23. Kelly DW, Dye C. Pheromones, kairomones and the aggregation dynamics of the sandfly *Lutzomyia longipalpis*. Anim Behav. 1997 Apr;53:721-31.
24. Alexander B. Sampling methods for phlebotomine sand flies. Med Vet Entomol. 2000 Jun;14(2):109-22.
25. Abbasi I, Cunio R, Warburg A. Identification of blood meals imbibed by phlebotomine sand flies using cytochrome b PCR and reverse line blotting. Vector-Borne and Zoonotic Diseases. 2009 Feb;9(1):79-86.
26. Bongiorno G, Habluetzel A, Khoury C, Maroli M. Host preferences of phlebotomine sand flies at a hypoendemic focus of canine leishmaniasis in central Italy. Acta Trop. 2003 Oct;88(2):109-16.
27. Pérez-Cutillas P, Muñoz C, Martínez-de la Puente J, Figuerola J, Navarro R, Ortuño M, et al. A spatial ecology study in a high diversity host community to understand blood feeding behavior of *Phlebotomus* sandfly vectors of *Leishmania*. Medical and Veterinary Entomology. 2020; In press.

28. Sallum MAM, Conn JE, Bergo ES, Laporta GZ, Chaves LSM, Bickersmith SA, et al. Vector competence, vectorial capacity of *Nyssorhynchus darlingi* and the basic reproduction number of *Plasmodium vivax* in agricultural settlements in the Amazonian Region of Brazil. *Malar J.* 2019 Apr 4;18.
29. Killick-Kendrick R. Phlebotomine vectors of the leishmaniasis - a review. *Med Vet Entomol.* 1990 Jan;4(1):1-24.
30. Ready PD. Biology of phlebotomine sand flies as vectors of disease agents. *Annual Review of Entomology*, Vol 58. 2013 2013;58:227-50.
31. Rioux JA, Aboulker JP, Lanotte G, Killick-Kendrick R, Martinidumas A. Ecology of Leishmaniasis in Southern France. 21. Influence of temperature on the development of *Leishmania infantum* - Nicolle, 1908 in *Phlebotomus - ariasi* Tommoir, 1921 - Experimental study. *Ann Parasitol Hum Comp.* 1985 1985;60(3):221-9.
32. Giorgobiani E, Lawyer PG, Babuadze G, Dolidze N, Jochim RC, Tskhvaradze L, et al. Incrimination of *Phlebotomus kandelakii* and *Phlebotomus balcanicus* as vectors of *Leishmania infantum* in Tbilisi, Georgia. *PLoS Negl Trop Dis.* 2012 Apr;6(4).
33. Doha S, Shehata MG. *Leishmania infantum*, MON-98 isolated from naturally infected *Phlebotomus langeroni* (Diptera, Psychodidae) in El Agamy, Egypt. *J Med Entomol.* 1992 Sep;29(5):891-3.
34. Leger N, Gramiccia M, Gradoni L, Maduloleblond G, Pesson B, Ferte H, et al. Isolation and typing of *Leishmania infantum* from *Phlebotomus neglectus* on the island of Corfu, Greece. *Trans R Soc Trop Med Hyg.* 1988 May-Jun;82(3):419-20.
35. Velo E, Bongiorno G, Kadriaj P, Myrseli T, Crilly J, Lika A, et al. The current status of phlebotomine sand flies in Albania and incrimination of *Phlebotomus neglectus* (Diptera, Psychodidae) as the main vector of *Leishmania infantum*. *PLoS One.* 2017 Jun 19;12(6).
36. Maroli M, Gramiccia M, Gradoni L. Natural infection of *Phlebotomus perfilliewi* with *Leishmania infantum* in a cutaneous leishmaniasis focus in The Abruzzi region, Italy. *Trans R Soc Trop Med Hyg.* 1987 1987;81(4):596-8.
37. Bettini S, Gramiccia M, Gradoni L, Atzeni MC. Leishmaniasis in Sardinia. 2. Natural infection of *Phlebotomus perniciosus* Newstead, 1911, by *Leishmania infantum* Nicolle, 1908, in the province of Cagliari. *Trans R Soc Trop Med Hyg.* 1986 1986;80(3):458-9.
38. Gonzalez E, Alvarez A, Ruiz S, Molina R, Jimenez M. Detection of high *Leishmania infantum* loads in *Phlebotomus perniciosus* captured in the leishmaniasis focus of southwestern Madrid region (Spain) by real time PCR. *Acta Trop.* 2017 Jul;171:68-73.
39. Leger N, Depaquit J, Ferte H, Rioux JA, Gantier JC, Gramiccia M, et al. Phlebotomine sand flies (Diptera psychodidae) of Cyprus. II - Isolation and typing of *Leishmania (Leishmania) infantum* Nicolle, 1908 (zymodeme MON 1) from *Phlebotomus (Larrousius) tobbi* Adler and Theodor, 1930. *Parasite.* 2000 Jun;7(2):143-6.
40. Svobodova M, Alten B, Zidkova L, Dvorak V, Hlavackova J, Myskova J, et al. Cutaneous leishmaniasis caused by *Leishmania infantum* transmitted by *Phlebotomus tobbi*. *Int J Parasitol.* 2009 Jan 15;39(2):251-6.
41. Lewis DJ, Ward RD. Transmission and vectors. In: Peters W, Killick-Kendrick R, editors. *The leishmaniasis in biology and medicine.* 1. London: Academic Press; 1987. p. 235-62.
42. Pesson B, Leger N, Maduloleblond G. Leishmaniasis in Greece -The sand flies of Ionian and Aegean islands. *Ann Parasitol Hum Comp.* 1984 1984;59(3):277-96.
43. Pehu B. Sur un cas de kala-azar survenu chez un enfant de 5 ans n'ayant jamais quitté les Vosges. *Lyon, France: Lyon Med;* 1931. p. 731-4.
44. Azizi K, Rassi Y, Javadian E, Motazedian MH, Asgari Q, Yaghoobi-Ershadi MR. First detection of *Leishmania infantum* in *Phlebotomus (Larrousius) major* (Diptera: Psychodidae) from Iran. *J Med Entomol.* 2008 Jul;45(4):726-31.
45. Emami MM, Yazdi M. Entomological survey of phlebotomine sand flies (Diptera: Psychodidae) in a focus of visceral leishmaniasis in central Iran. *J Vector Borne Dis.* 2008 2008-Mar;45(1):38-43.
46. Maroli M, Felicangeli MD, Bichaud L, Charrel RN, Gradoni L. Phlebotomine sand flies and the spreading of leishmaniasis and other diseases of public health concern. *Med Vet Entomol.* 2013 Jun;27(2):123-47.
47. Boubidi SC, Benallal K, Boudrissa A, Bouiba L, Bouchareb B, Garni R, et al. *Phlebotomus sergenti* (Parrot, 1917) identified as *Leishmania killicki* host in Ghardaia, south Algeria. *Microb Infect.* 2011 Jul;13(7):691-6.
48. Tabbabi A, Bousslimi N, Rhim A, Aoun K, Bouratbine A. Short report: first report on natural infection of *Phlebotomus sergenti* with *Leishmania* promastigotes in the cutaneous leishmaniasis focus in south-eastern Tunisia. *Am J Trop Med Hyg.* 2011 Oct;85(4):646-7.
49. Depaquit J, Ferte H, Leger N, Lefranc F, Alves-Pires C, Hanafi H, et al. ITS 2 sequences heterogeneity in *Phlebotomus sergenti* and *Phlebotomus similis* (Diptera, Psychodidae): possible consequences in their ability to transmit *Leishmania tropica*. *Int J Parasitol.* 2002 Aug;32(9):1123-31.
50. Christodoulou V, Antoniou M, Ntais P, Messaritakis I, Iovic V, Dedet J-P, et al. Re-emergence of visceral and cutaneous leishmaniasis in the Greek island of Crete. *Vector-Borne and Zoonotic Diseases.* 2012 Mar;12(3):214-22.
51. Volf P, Myskova J. Sand flies and *Leishmania*: specific versus permissive vectors. *Trends Parasitol.* 2007 Mar;23(3):91-2.
52. Dvorak V, Shaw J, Volf P. Parasite biology: the vector. In: Bruchi F, Gradoni L, editors. *The Leishmaniasis: old neglected tropical diseases.* Cham, Switzerland: Springer; 2018. p. 31-76.
53. Alkan C, Bichaud L, de Lamballerie X, Alten B, Gould EA, Charrel RN. Sandfly-borne phleboviruses of Eurasia and Africa: epidemiology, genetic diversity, geographic range, control measures. *Antiviral Res.* 2013 Oct;100(1):54-74.
54. Mihalca AD, Cazan CD, Sulesco T, Dumitrache MO. A historical review on vector distribution and epidemiology of human and animal leishmaniasis in Eastern Europe. *Res Vet Sci.* 2019 Apr;123:185-91.

55. Maroli M, Rossi L, Baldelli R, Capelli G, Ferroglio E, Genchi C, et al. The northward spread of leishmaniasis in Italy: evidence from retrospective and ongoing studies on the canine reservoir and phlebotomine vectors. *Trop Med Int Health*. 2008 Feb;13(2):256-64.
56. Dereure J, Vanwambeke SO, Male P, Martinez S, Pratlong F, Balard Y, et al. The potential effects of global warming on changes in canine leishmaniasis in a focus outside the classical area of the disease in Southern France. *Vector-Borne and Zoonotic Diseases*. 2009 Dec;9(6):687-94.
57. Baldelli R, Battelli G, Maroli M, Mollicone E, Gudi A, Stegagno G, et al. A new stable focus of canine leishmaniasis in northern Italy. *Parassitologia*. 2001 2001-Dec;43(4):151-3.
58. Galvez R, Descalzo MA, Guerrero I, Miro G, Molina R. Mapping the current distribution and predicted spread of the leishmaniasis sand fly vector in the Madrid region (Spain) based on environmental variables and expected climate change. *Vector-Borne and Zoonotic Diseases*. 2011 Jul;11(7):799-806.
59. Martin Sanchez J, Morales-Yuste M, Acedo-Sanchez C, Baron S, Diaz V, Morillas-Marquez F. Canine leishmaniasis in south-eastern Spain. *Emerg Infect Dis*. 2009 May;15(5):795-8.
60. Shaw SE, Langton DA, Hillman TJ. Canine leishmaniasis in the United Kingdom: a zoonotic disease waiting for a vector? *Vet Parasitol*. 2009 Aug 26;163(4):281-5.
61. Menn B, Lorentz S, Naucke TJ. Imported and travelling dogs as carriers of canine vector-borne pathogens in Germany. *Parasites & Vectors*. 2010 Apr 8;3.
62. Otranto D, Dantas-Torres F, Mihalca AD, Traub RJ, Lappin M, Baneth G. Zoonotic parasites of sheltered and stray dogs in the era of the global economic and political crisis. *Trends Parasitol*. 2017 Oct;33(10):813-25.
63. Svobodova V, Svoboda M, Friedlaenderova L, Drahotsky P, Bohacova E, Baneth G. Canine leishmaniasis in three consecutive generations of dogs in Czech Republic. *Vet Parasitol*. 2017 Apr 15;237:122-4.
64. Toepp AJ, Schaut RG, Scott BD, Mathur D, Berens AJ, Petersen CA. *Leishmania* incidence and prevalence in U.S. hunting hounds maintained via vertical transmission. *Veterinary Parasitology, regional studies and reports*. 2017 2017-12;10:75-81.
65. Ravel C, Cortes S, Pratlong F, Morio F, Dedet J-P, Campino L. First report of genetic hybrids between two very divergent *Leishmania* species: *Leishmania infantum* and *Leishmania major*. *Int J Parasitol*. 2006 Nov;36(13):1383-8.
66. Volf P, Benkova I, Myskova J, Sadlova J, Campino L, Ravel C. Increased transmission potential of *Leishmania major*/*Leishmania infantum* hybrids. *Int J Parasitol*. 2007 May;37(6):589-93.
67. Campino L, Cortes S, Dionisio L, Neto L, Afonso MO, Maia C. The first detection of *Leishmania major* in naturally infected *Sergentomyia minuta* in Portugal. *Mem Inst Oswaldo Cruz*. 2013 Jun;108(4):516-8.
68. Bongiorno G, Di Muccio T, Bianchi R, Gramiccia M, Gradoni L. Laboratory transmission of an Asian strain of *Leishmania tropica* by the bite of the southern European sand fly *Phlebotomus perniciosus*. *Int J Parasitol*. 2019 May;49(6):417-21.
69. Antoniou M, Gramiccia M, Molina R, Dvorak V, Volf P. The role of indigenous phlebotomine sand flies and mammals in the spreading of leishmaniasis agents in the Mediterranean region. *Eurosurveillance*. 2013 Jul 25;18(30):54-61.
70. Ozbilgin A, Harman M, Karakus M, Bart A, Toz S, Kurt O, et al. Leishmaniasis in Turkey: visceral and cutaneous leishmaniasis caused by *Leishmania donovani* in Turkey. *Acta Trop*. 2017 Sep;173:90-6.
71. Seblova V, Myskova J, Hlavacova J, Votypka J, Antoniou M, Volf P. Natural hybrid of *Leishmania infantum*/*L. donovani*: development in *Phlebotomus tobbi*, *P. perniciosus* and *Lutzomyia longipalpis* and comparison with non-hybrid strains differing in tissue tropism. *Parasites & Vectors*. 2015 Nov 25;8.
72. Gonzalez E, Jimenez M, Hernandez S, Martin-Martin I, Molina R. Phlebotomine sand fly survey in the focus of leishmaniasis in Madrid, Spain (2012-2014): seasonal dynamics, *Leishmania infantum* infection rates and blood meal preferences. *Parasites & Vectors*. 2017 Aug 1;10.
73. Tomassone L, Berriatua E, De Sousa R, Duscher GG, Mihalca AD, Silaghi C, et al. Neglected vector-borne zoonoses in Europe: into the wild. *Vet Parasitol*. 2018 Feb 15;251:17-26.
74. Risueno J, Ortuno M, Perez-Cutillas P, Goyena E, Maia C, Cortes S, et al. Epidemiological and genetic studies suggest a common *Leishmania infantum* transmission cycle in wildlife, dogs and humans associated to vector abundance in south-east Spain. *Vet Parasitol*. 2018 Aug 15;259:61-7.
75. Alonso F, Gimenez Font P, Manchon M, Ruiz de Ybanez R, Segovia M, Berriatua E. Geographical variation and factors associated to seroprevalence of canine leishmaniasis in an endemic Mediterranean area. *Zoonoses and Public Health*. 2010 Aug;57(5):318-28.
76. Sanchez CA, Sanchez JM, Bernal IDV, Marin MCS, Louassini M, Maldonado JA, et al. Leishmaniasis eco-epidemiology in the Alpujarra region (Granada province, southern Spain). *Int J Parasitol*. 1996 Mar;26(3):303-10.
77. Gebre-Michael T, Malone JB, Balkew M, Ali A, Berhe N, Hailu A, et al. Mapping the potential distribution of *Phlebotomus martini* and *P. orientalis* (Diptera: Psychodidae), vectors of kala-azar in East Africa by use of geographic information systems. *Acta Trop*. 2004 Mar;90(1):73-86.
78. Franco AO, Davies CR, Mylne A, Dedet J-P, Gallego M, Ballart C, et al. Predicting the distribution of canine leishmaniasis in western Europe based on environmental variables. *Parasitology*. 2011 Dec;138(14):1878-91.
79. Perez-Cutillas P, Goyena E, Chitimia L, De la Rúa P, Bernal LJ, Fisa R, et al. Spatial distribution of human asymptomatic *Leishmania infantum* infection in southeast Spain: a study of environmental, demographic and social risk factors. *Acta Trop*. 2015 Jun;146:127-34.

80. Risueno J, Munoz C, Perez-Cutillas P, Goyena E, Gonzalez M, Ortuno M, et al. Understanding *Phlebotomus perniciosus* abundance in south-east Spain: assessing the role of environmental and anthropic factors. *Parasites & Vectors*. 2017 Apr 19;10.
81. Arce A, Estirado A, Ordobas M, Sevilla S, Garcia N, Moratilla L, et al. Re-emergence of leishmaniasis in Spain: community outbreak in Madrid, Spain, 2009 to 2012. *Eurosurveillance*. 2013 Jul 25;18(30):5-13.
82. Gradoni L. A brief introduction to leishmaniasis epidemiology. In: Bruchi F, Gradoni L, editors. *The Leishmaniasis: old neglected tropical diseases*. Cham, Switzerland: Springer; 2018. p. 1-14.
83. Gramiccia M, Scalone A, Di Muccio T, Orsini S, Fiorentino E, Gradoni L. The burden of visceral leishmaniasis in Italy from 1982 to 2012: a retrospective analysis of the multi-annual epidemic that occurred from 1989 to 2009. *Eurosurveillance : bulletin European sur les maladies transmissibles = European communicable disease bulletin*. 2013 2013 Jul;18(29):20535.
84. Alvar J, Canavate C, Gutierrez Solar B, Jimenez M, Laguna F, Lopez Velez R, et al. Leishmania and human immunodeficiency virus coinfection: the first 10 years. *Clin Microbiol Rev*. 1997 Apr;10(2):298-+.
85. Botet Fregola J, Portus Vinyeta M. Leishmaniasis in peninsular Spain. A historical-bibliographic review (1912-1985). *Rev Sanid Hig Publica (Madr)*. 1993;67(4):255-66.
86. World Health Organization. Report of the Interregional meeting on leishmaniasis among neighbouring endemic countries in the Eastern Mediterranean, African and European regions, Amman, Jordan, 23–25 September 2018. . WHO, 2019 Contract No.: WHO-EM/CTD/081/E.
87. Otranto D, Dantas-Torres F. The prevention of canine leishmaniasis and its impact on public health. *Trends Parasitol*. 2013 Jul;29(7):339-45.
88. Galvez R, Montoya A, Fontal F, Martinez De Murguia L, Miro G. Controlling phlebotomine sand flies to prevent canine *Leishmania infantum* infection: a case of knowing your enemy. *Res Vet Sci*. 2018 Dec;121:94-103.
89. Maroli M, Mizioni V, Siragusa C, D'Orazi A, Gradoni L. Evidence for an impact on the incidence of canine leishmaniasis by the mass use of deltamethrin-impregnated dog collars in southern Italy. *Med Vet Entomol*. 2001 Dec;15(4):358-63.
90. Gavgani ASM, Hodjati MH, Mohite H, Davies CR. Effect of insecticide-impregnated dog collars on incidence of zoonotic visceral leishmaniasis in Iranian children: a matched-cluster randomised trial. *Lancet*. 2002 Aug 3;360(9330):374-9.
91. Manzillo VF, Oliva G, Pagano A, Manna L, Maroli M, Gradoni L. Deltamethrin-impregnated collars for the control of canine leishmaniasis: evaluation of the protective effect and influence on the clinical outcome of *Leishmania* infection in kennelled stray dogs. *Vet Parasitol*. 2006 Nov 30;142(1-2):142-5.
92. Otranto D, Paradies P, Lia RP, Latrofa MS, Testini G, Cantacessi C, et al. Efficacy of a combination of 10% imidacloprid/50% permethrin for the prevention of leishmaniasis in kennelled dogs in an endemic area. *Vet Parasitol*. 2007 Mar 31;144(3-4):270-8.
93. Otranto D, de Caprariis D, Lia RP, Tarallo V, Lorusso V, Testini G, et al. Prevention of endemic canine vector-borne diseases using imidacloprid 10% and permethrin 50% in young dogs: a longitudinal field study. *Vet Parasitol*. 2010 Sep 20;172(3-4):323-32.
94. Ferroglio E, Poggi M, Trisciuglio A. Evaluation of 65% permethrin spot-on and deltamethrin-impregnated collars for canine *Leishmania infantum* infection prevention. *Zoonoses and Public Health*. 2008 Apr;55(3):145-8.
95. Courtenay O, Bazmani A, Parvizi P, Ready PD, Cameron MM. Insecticide-impregnated dog collars reduce infantile clinical visceral leishmaniasis under operational conditions in NW Iran: a community-wide cluster randomised trial. *PLoS Negl Trop Dis*. 2019 Mar;13(3).
96. Reithinger R, Coleman PG, Alexander B, Vieira EP, Assis G, Davies CR. Are insecticide-impregnated dog collars a feasible alternative to dog culling as a strategy for controlling canine visceral leishmaniasis in Brazil? *Int J Parasitol*. 2004 Jan;34(1):55-62.
97. Goyena E, Perez-Cutillas P, Chitimia L, Risueno J, Garcia-Martinez JD, Bernal LJ, et al. A cross-sectional study of the impact of regular use of insecticides in dogs on canine leishmaniasis seroprevalence in southeast Spain. *Prev Vet Med*. 2016 Feb 1;124:78-84.
98. Killick-Kendrick R. The biology and control of phlebotomine sand flies. *Clin Dermatol*. 1999 May-Jun;17(3):279-89.
99. Alexander B, Maroli M. Control of phlebotomine sand flies. *Med Vet Entomol*. 2003 Mar;17(1):1-18.
100. Iriso A, Tello A, González-Mora D, Vázquez M, Molina R, Jiménez M, et al. Control del Vector. In: Madrid DGdSPCdS-Cd, editor. *Brotos de leishmaniasis en Fuenlabrada y otros municipios de la Comunidad de Madrid El papel de las liebres y los conejos como reservorios*. 1. Madrid: Biblioteca Virtual de la Comunidad de Madrid; 2017. p. 177-90.
101. Vilas F, Fúster F. Gestión del brote de Leishmaniosis. In: Madrid DGdSPCdS-Cd, editor. *Brotos de leishmaniasis en Fuenlabrada y otros municipios de la Comunidad de Madrid El papel de las liebres y los conejos como reservorios*. 1. Madrid: Biblioteca Virtual de la Comunidad de Madrid; 2017. p. 55-63.
102. Tello A, González-Mora D, Outerelo R, Iriso A, Vázquez M. Los flebotomos del brote de leishmaniasis en el suroeste de la Comunidad de Madrid (Diptera, Psychodidae, Phlebotominae). *Boletín Real Sociedad Española de Historia Natural Sección Biológica [Internet]*. 2015; 109:57-64.
103. Vázquez A, Tello A, Iriso A, González D, Outerelo R, Martínez M, et al. Vigilancia del vector. In: Madrid DGdSPCdS-Cd, editor. *Brotos de leishmaniasis en Fuenlabrada y otros municipios de la Comunidad de Madrid El papel de las liebres y los conejos como reservorios*. 1: Biblioteca Virtual de la Comunidad de Madrid; 2017. p. 171-6.
104. Olmedo Luceron C, Perez Meixeira AM, Iriso Calle A, Aranguez Ruiz E, Abad Sanz I. Evolution and spatial analysis of leishmaniasis cases in a rural area bordering a community outbreak in Madrid: 2001-2017. *Rev Esp Salud Publica*. 2018 Nov;92.

105. Nery Costa CH. How effective is dog culling in controlling zoonotic visceral leishmaniasis? A critical evaluation of the science, politics and ethics behind this public health policy. *Rev Soc Bras Med Trop.* 2011 Mar-Apr;44(2):232-42.
106. de Sousa-Paula LC, da Silva LG, da Silva Sales KG, Dantas-Torres F. Failure of the dog culling strategy in controlling human visceral leishmaniasis in Brazil: a screening coverage issue? *PLoS Negl Trop Dis.* 2019 Jun;13(6).
107. Chaniotis BN, Parsons RE, Harlan HJ, Correa MA. A pilot study to control Phlebotomine sand flies (Diptera, Psychodidae) in a neotropical rain-forest. *J Med Entomol.* 1982 1982;19(1):1-5.
108. Ready PD, Arias JR, Freitas RA. A pilot study to control *Lutzomyia umbratilis* (Diptera:Psychodidae), the major vector of *Leishmania braziliensis guyanensis*, in a peri-urban rainforest of Manaus, Amazonas State, Brazil. *Mem Inst Oswaldo Cruz.* 1985 1985;80(1):27-36.
109. Robert LL, Perich MJ. Phlebotomine sand fly (Diptera, Psychodidae) control using a residual pyrethroid insecticide. *J Am Mosq Control Assoc.* 1995 Jun;11(2):195-9.
110. Tesh RB, Papaevangelou G. Effect of insecticide spraying for malaria control on incidence of sandfly fever in Athens, Greece. *Am J Trop Med Hyg.* 1977 1977;26(1):163-6.
111. Sharma U, Singh S. RETRACTED: Insect vectors of *Leishmania*: distribution, physiology and their control. *J Vector Borne Dis.* 2008 2008-Dec;45(4):255-72.
112. Poche DM, Torres-Poche Z, Yeszhanov A, Poche RM, Belyaev A, Dvorak V, et al. Field evaluation of a 0.005% fipronil bait, orally administered to *Rhombomys opimus*, for control of fleas (Siphonaptera: Pulicidae) and phlebotomine sand flies (Diptera: Psychodidae) in the Central Asian Republic of Kazakhstan. *PLoS Negl Trop Dis.* 2018 Jul;12(7).
113. Warburg A, Faiman R. Research priorities for the control of phlebotomine sand flies. *J Vector Ecol.* 2011 Mar;36:S10-S6.
114. Mueller GC, Schlein Y. Different methods of using attractive sugar baits (ATSB) for the control of *Phlebotomus papatasi*. *J Vector Ecol.* 2011 Mar;36:S64-S70.
115. Qualls WA, Mueller GC, Khallaayoune K, Revay EE, Zhioua E, Kravchenko VD, et al. Control of sand flies with attractive toxic sugar baits (ATSB) and potential impact on non-target organisms in Morocco. *Parasites & Vectors.* 2015 Feb 8;8.
116. Saghafipour A, Vatandoost H, Zahraei-Ramazani AR, Yaghoobi-Ershadi MR, Rassi Y, Jooshin MK, et al. Control of zoonotic cutaneous leishmaniasis vector, *Phlebotomus papatasi*, using attractive toxic sugar baits (ATSB). *PLoS One.* 2017 Apr 20;12(4).
117. McDermott EG, Morris EK, Garver LS. Sodium ascorbate as a potential toxicant in attractive sugar baits for control of adult mosquitoes (Diptera: Culicidae) and sand flies (Diptera: Psychodidae). *J Med Entomol.* 2019 Sep;56(5):1359-67.
118. Robert LL, Perich MJ, Schlein Y, Jacobson RL, Wirtz RA, Lawyer PG, et al. Phlebotomine sand fly control using bait-fed adults to carry the larvicide *Bacillus sphaericus* to the larval habitat. *J Am Mosq Control Assoc.* 1997 Jun;13(2):140-4.
119. Robert LL, Perich MJ, Schlein Y, Jacobson JL. *Bacillus sphaericus* inhibits hatching of phlebotomine sand fly eggs. *J Am Mosq Control Assoc.* 1998 Sep;14(3):351-2.
120. Wahba MM. The influence of *Bacillus sphaericus* on the biology and histology of *Phlebotomus papatasi*. *J Egypt Soc Parasitol.* 2000 2000-Apr;30(1):315-23.
121. Emami MM, Yazdi M, Guillet P. Efficacy of Olyset long-lasting bednets to control transmission of cutaneous leishmaniasis in Iran. *Eastern Mediterranean health journal = La revue de sante de la Mediterranee orientale = al-Majallah al-sihhiyah li-sharq al-mutawassit.* 2009;15(5):1075-83.
122. Gunay F, Karakus M, Oguz G, Dogan M, Karakaya Y, Ergan G, et al. Evaluation of the efficacy of Olyset (R) Plus in a village-based cohort study in the Cukurova Plain, Turkey, in an area of hyperendemic cutaneous leishmaniasis. *J Vector Ecol.* 2014 Dec;39(2):395-405.
123. Chowdhury R, Chowdhury V, Faria S, Akter S, Dash AP, Bhattacharya SK, et al. Effect of insecticide-treated bed nets on visceral leishmaniasis incidence in Bangladesh. A retrospective cohort analysis. *PLoS Negl Trop Dis.* 2019 Sep;13(9).
124. Picado A, Ostyn B, Rijal S, Sundar S, Singh P, Chappuis F, et al. Long-lasting insecticidal nets to prevent visceral leishmaniasis in the Indian subcontinent; methodological lessons learned from a cluster randomised controlled trial. *PLoS Negl Trop Dis.* 2015 Apr;9(4).
125. Nyunt MH, Aye KM, Kyaw MP, Kyaw TT, Hlaing T, Oo K, et al. Challenges in universal coverage and utilization of insecticide-treated bed nets in migrant plantation workers in Myanmar. *Malar J.* 2014 Jun 2;13.
126. Faraj C, Yukich J, Adlaoui EB, Wahabi R, Mnzava AP, Kaddaf M, et al. Effectiveness and cost of insecticide-treated bed nets and indoor residual spraying for the control of cutaneous leishmaniasis: a cluster-randomized control trial in Morocco. *Am J Trop Med Hyg.* 2016 Mar;94(3):679-85.
127. Boelaert M, Burza S, Romero G. Control and public health aspects. In: Bruchi F, Gradoni L, editors. *The Leishmaniasis: old neglected tropical diseases.* Cham, Switzerland: Springer; 2018. p. 227-45.

Table 1. Presence (1) and absence (blank) of confirmed and suspected *Leishmania* spp. vectors in Europe and neighbouring countries and regions.

Country/region	<i>Phlebotomus</i> species															
	<i>alexandri</i>	<i>ariasi</i>	<i>balcanicus</i>	<i>halepensis</i>	<i>kandelakii</i> s.l.	<i>langeroni</i>	<i>longicuspis</i>	<i>major</i> s.l.	<i>mascittii</i>	<i>papatasi</i>	<i>perfiliewi</i> s.l.	<i>pernicius</i>	<i>sergenti</i>	<i>simici</i>	<i>similis</i>	<i>tobbi</i>
Canary Islands		1											1			
Germany								1	1			1				
Spain (continental)	1	1				1	1		1	1		1	1			
Balearic Islands										1		1	1			
Switzerland								1	1	1		1				
France (continental)		1							1	1		1	1			
Portugal		1								1		1	1			
Belgium									1							
Corsica									1	1		1	1			
Morocco	1	1				1	1			1	1	1	1			
Tunisia	1	1		1		1	1			1	1	1	1			
Azerbaijan	1		1	1	1					1	1	1	1		1	1
Iran	1		1		1			1		1	1		1		1	1
Iraq	1									1	1		1			1
Libya	1					1	1			1		1	1			1
Algeria	1	1				1	1		1	1	1	1	1			
Israel	1			1					1	1	1		1	1		1
Palestine	1			1		1		1	1	1	1		1			1
Turkey (Anatolia)	1		1	1	1			1	1	1	1		1	1	1	1
Armenia	1		1		1			1		1	1		1			1
Georgia			1	1	1			1		1	1		1	1		
Cyprus Island	1							1	1	1	1	1	1			1
Kuwait										1			1			
Syria	1			1				1		1	1		1			1
Ukraine	1		1					1		1	1				1	
Egypt	1					1		1		1			1			
Jordan	1			1				1	1	1	1		1			1
Lebanon	1		1		1	1		1		1			1	1		1
Russia (south-western)					1			1		1			1	1		
Moldova										1	1					
Crimean Peninsula	1		1					1		1	1	1			1	
Hungary								1	1	1	1					
Bulgaria	1				1			1		1	1		1	1		1

Country/region	Phlebotomus species															
	<i>alexandri</i>	<i>ariasi</i>	<i>balcanicus</i>	<i>halepensis</i>	<i>kandelakii</i> s.l.	<i>langeroni</i>	<i>longicuspis</i>	<i>major</i> s.l.	<i>mascittii</i>	<i>papatasi</i>	<i>perfiliewi</i> s.l.	<i>perniciosus</i>	<i>sergenti</i>	<i>simici</i>	<i>similis</i>	<i>tobbi</i>
Croatia							1	1	1	1	1					1
Italy (continental)		1					1	1	1	1	1	1				1
Kosovo*	1						1		1	1				1		1
Malta							1		1	1	1	1				
Sicily							1	1	1	1	1	1				
Bosnia & Herzegovina			1				1	1	1					1		1
Slovakia								1								
Crete	1						1	1	1	1				1	1	1
Sardinia									1	1						
Romania	1		1				1		1	1		1				
Greece	1		1		1		1	1	1	1		1	1	1	1	1
North Macedonia	1		1				1	1	1	1	1	1	1	1	1	1
Montenegro			1		1		1		1	1	1		1			1
Serbia	1		1				1	1	1	1	1	1	1			1
Slovenia							1	1	1		1					
Albania							1		1	1					1	1
Austria								1								
Turkey (Thrace)							1			1						1

	<i>validated</i>
	<i>no reference</i>
	<i>unpublished VectorNet data</i>

* This designation is without prejudice to positions on status, and is in line with UNSCR 1244 and the ICJ Opinion on the Kosovo Declaration of Independence.