Tuberculosis surveillance and monitoring in Europe 2017



Poland

Total population at 10th August 2016 by EUROSTAT: 38 005 614

Tuberculosis case notifications, 2015

| Total number of cases | 6430 | | |
|--|---------|------------|--|
| Notification rate per100 000 | 16 | 16.9 | |
| New* & relapses | 6 2 3 7 | | |
| New* & relapses notification rate per100 000 | 16 | 16.4 | |
| Pulmonary | 6078 | (94.5%) | |
| of which microscopy-positive | 2784 | (45.8%) | |
| of which laboratory confirmed | 4472 | (73.6%) | |
| Laboratory-confirmed TB cases | 4630 | (72.0%) | |
| Mean age of new native TB cases | 53.7 | 53.7 years | |
| Mean age of new foreign TB cases | 31.0 | 31.0 years | |
| Foreign origin of all TB cases | 52 | (0.8%) | |
| New (not previously treated) | 5757 | (89.5%) | |
| | | | |

^{*} Cases with unknown previous treatment included in new cases.

Drug resistance surveillance & TB-HIV co-infection, 2015

| Completeness of DRS data* | Υ | es |
|------------------------------------|----------|---------|
| Completeness of HIV data** | Υ | es |
| Case-linked data reporting | Yes | |
| Cases with DST results | 4242 | (91.6%) |
| Estimated MDR N, (best-low-high) | 60-42-79 | |
| Pulmonary MDR TB cases notified | 32 | (0.8%) |
| of which XDR TB cases | 4 | (17.4%) |
| Notified MDR TB | 35 | (0.8%) |
| of which XDR TB cases | 4 | (16.7%) |
| TB cases tested for HIV | - | - |
| HIV-positive TB cases | - | - |
| of these on antiretroviral therapy | - | - |
| | | |

^{*} National coverage 100% or culturing 90%, C+/All TB cases 50%, DST done for C+75%, EQA 95%.


** More than 50% of TB cases tested for HIV.

Treatment outcome monitoring


| Geographical coverage | Nat | ional | | |
|----------------------------|---|---------|--|---------|
| Outcome cohort | New culture positive pulmonary TB cases notified in 2014* | | All MDR TB cases notified in 2013** | |
| Case-linked data reporting | ١ | 'es | | |
| Cases notified | 4153 | | 40 | |
| Success | 2434 | (58.6%) | 6 | (15.0%) |
| Died | 392 | (9.4%) | 8 | (20.0%) |
| Failed | 5 | (0.1%) | 0 | (0.0%) |
| Lost to follow-up | 247 | (5.9%) | 7 | (17.5%) |
| Still on treatment | 0 | (0.0%) | 2 | (5.0%) |
| Not evaluated | 1075 | (25.9%) | 17 | (42.5%) |

^{*} Treatment outcome as presented is treatment outcome after 12 months.


Tuberculosis notification rates by treatment history, 2006-2015



New and relapsed TB cases - notification rates by age group, 2006-2015




Tuberculosis cases by geographical origin, 2006-2015




TB-HIV co-infection, 2006-2015

Data not available

MDR TB cases by previous treatment history, 2006-2015



Treatment outcome, new culture-confirmed pulmonary TB cases,



Source: European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Tuberculosis surveillance and monitoring in Europe 2017. Stockholm: ECDC; 2017

^{**} Treatment outcome as presented is treatment outcome after 24 months.