

SURVEILLANCE REPORT

Weekly influenza surveillance overview

12 March 2010

Main surveillance developments in week 09/2010 (01 Mar 2010 – 07 Mar 2010)

This first page contains the main developments this week and can be printed separately or together with the more detailed information following.

- All 25 reporting countries reported low intensity for week 09/2010.
- Regional geographic spread was reported in Austria and Greece, local spread in Italy and Malta, and all other countries reported sporadic or no activity.
- Of the 425 specimens collected by sentinel physicians, 20 (4.7%) were positive for influenza virus, mainly the pandemic virus, but there is some circulation of influenza B viruses; this is consistent with a declining trend.
- In cases of influenza-like illness seen by sentinel physicians, less than 5% are now positive for influenza virus.
- The number of reported SARI cases continued to decline. Only five SARI cases were reported during week 09/2010.
- Even though globally the world remains in pandemic phase 6, influenza activity caused by the 2009 pandemic influenza A(H1N1) virus is well past its winter peak in EU/EEA countries.

Sentinel surveillance of influenza like-illness (ILI)/ acute respiratory illness (ARI): All 25 reporting countries reported low ILI/ARI activity. For the trend indicator, 24 countries reported a stable or declining trend, with only Lithuania reporting an increasing trend. For more information, [click here](#).

Virological surveillance: Sentinel physicians collected 425 respiratory specimens, 20 (4.7%) of which were positive for influenza virus. Since week 40/2009, 99% of the viruses detected in sentinel specimens were 2009 pandemic influenza A(H1N1) virus. For more information, [click here](#).

Aggregate numbers of 2009 pandemic influenza (H1N1) deaths: During week 09/2010, six countries reported six deaths. For more information, [click here](#).

Hospital surveillance of severe acute respiratory infection (SARI): During week 09/2010, five SARI cases were reported. For more information, [click here](#).

Qualitative reporting: For more information, [click here](#).

Sentinel surveillance (ILI/ARI)

Weekly analysis – epidemiology


In week 9/2010, 25 out of 29 countries reported epidemiological data. For the activity intensity indicator, all countries reported low intensity. For the trend indicator, 24 countries reported stable or declining trend, with only Lithuania reporting increasing trend (Map 1 and Table 1).

For the geographic spread indicator, regional geographic spread was reported in Austria and Greece, local spread reported in Italy and Malta, and all other countries reported sporadic or no activity (Map 2 and Table 1).

Since week 03/2010, all countries have reported medium or low intensity activity, with the number of countries reporting medium intensity decreasing from five to zero in week 09, when all the countries reported low intensity.

For the majority of countries that reported age specific incidence of ILI and/or ARI, the most affected age group was 0–14 years.

Map 1: Intensity for week 09/2010



* A type/subtype is reported as dominant when > 40 % of all samples are positive for the type/subtype.


Legend:

Low	No influenza activity or influenza at baseline levels	-	Decreasing clinical activity
Medium	Usual levels of influenza activity	+	Increasing clinical activity
High	Higher than usual levels of influenza activity	=	Stable clinical activity
Very high	Particularly severe levels of influenza activity	A	Type A
		A(H1)v	Type A, Subtype H1v
		A(H1N1)v	Type A, Subtype H1N1v
		B	Type B

Map 2: Geographic spread for week 09/2010

Geographic spread

- No Report
- No Activity
- Sporadic
- Local
- Regional
- Widespread



- Liechtenstein
- Luxembourg
- Malta

(C) ECDC/Dundas/TESSy

* A type/subtype is reported as dominant when > 40 % of all samples are positive for the type/subtype.

Legend:

No activity	No evidence of influenza virus activity (clinical activity remains at baseline levels)	-	Decreasing clinical activity
Sporadic	Isolated cases of laboratory confirmed influenza infection	+	Increasing clinical activity
Local outbreak	Increased influenza activity in local areas (e.g. a city) within a region, or outbreaks in two or more institutions (e.g. schools) within a region (laboratory confirmed)	=	Stable clinical activity
Regional activity	Influenza activity above baseline levels in one or more regions with a population comprising less than 50% of the country's total population (laboratory confirmed)	A	Type A
Widespread	Influenza activity above baseline levels in one or more regions with a population comprising 50% or more of the country's population (laboratory confirmed)	A(H1)v	Type A, Subtype H1v
		A(H1N1)v	Type A, Subtype H1N1v
		B	Type B

Table 1: Epidemiological and virological overview by country

Country	Intensity	Geographic spread	Trend	No. of sentinel swabs	Dominant type	Percentage positive*	ILI per 100.000	ARI per 100.000	Epidemiological overview	Virological overview
Austria	Low	Regional	Stable	6	A(H1N1)v	0.0	-	20.2	Graphs	Graphs
Belgium	Low	Sporadic	Stable	-	-	-	65.5	1411.2	Graphs	Graphs
Bulgaria	Low	No activity	Decreasing	0	None	-	-	741.2	Graphs	Graphs
Cyprus				-	-	-	-	-	Graphs	Graphs
Czech Republic	Low	Sporadic	Stable	17	A	11.8	33.4	884.8	Graphs	Graphs
Denmark	Low	No activity	Stable	13	None	0.0	49.9	0.0	Graphs	Graphs
Estonia	Low	Sporadic	Decreasing	14	None	0.0	5.8	247.5	Graphs	Graphs
Finland				-	-	-	-	-	Graphs	Graphs
France	Low	Sporadic	Stable	35	None	0.0	-	1197.2	Graphs	Graphs
Germany	Low	Sporadic	Stable	28	None	10.7	-	1081.3	Graphs	Graphs
Greece	Low	Regional	Decreasing	3	A(H1N1)v	33.3	124.6	-	Graphs	Graphs
Hungary	Low	Sporadic	Decreasing	56	None	7.1	110.0	-	Graphs	Graphs
Iceland	Low	No activity	Stable	-	-	-	0.6	-	Graphs	Graphs
Ireland	Low	Sporadic	Stable	8	A(H1N1)v	0.0	8.8	-	Graphs	Graphs
Italy	Low	Local	Decreasing	34	A(H1N1)v	8.8	171.6	-	Graphs	Graphs
Latvia	Low	Sporadic	Decreasing	0	None	-	0.9	1144.8	Graphs	Graphs
Lithuania	Low	Sporadic	Increasing	-	-	-	2.3	566.3	Graphs	Graphs
Luxembourg				16	None	0.0	-*	-*	Graphs	Graphs
Malta	Low	Local	Stable	-	-	-	-*	-*	Graphs	Graphs
Netherlands	Low	Sporadic	Stable	8	None	0.0	31.7	-	Graphs	Graphs
Norway				0	None	-	-	-	Graphs	Graphs
Poland	Low	Sporadic	Decreasing	8	None	0.0	79.4	-	Graphs	Graphs
Portugal	Low	Sporadic	Stable	4	A(H1)v	0.0	9.9	-	Graphs	Graphs
Romania	Low	No activity	Stable	14	A(H1N1)v	14.3	0.5	793.7	Graphs	Graphs
Slovakia	Low	Sporadic	Decreasing	3	None	0.0	159.3	1383.5	Graphs	Graphs
Slovenia	Low	No activity	Stable	6	None	0.0	0.0	1050.0	Graphs	Graphs
Spain	Low	No activity	Stable	58	None	1.7	10.4	-	Graphs	Graphs
Sweden	Low	No activity	Stable	22	B	9.1	3.3	-	Graphs	Graphs
UK - England	Low	Sporadic	Stable	47	A(H1N1)v	3.6	6.3	379.7	Graphs	Graphs
UK - Northern Ireland	Low	No activity	Decreasing	4	None	0.0	18.5	398.3	Graphs	Graphs
UK - Scotland	Low	Sporadic	Stable	21	A(H1N1)v	4.8	3.0	209.0	Graphs	Graphs
UK - Wales				-	-	-	-	-	Graphs	Graphs
Europe				425		4.7				Graphs

*Incidence per 100 000 is not calculated for these countries as no population denominator is provided.

Note: Liechtenstein is not reporting to the European Influenza Surveillance Network

Description of the system

This surveillance is based on nationally organized sentinel networks of physicians, mostly general practitioners (GPs), covering at least 1–5% of the population in their countries. All EU/EEA Member States (except Liechtenstein) are participating. Depending on their country's choice, each sentinel physician reports the weekly number of patients seen with influenza-like illness (ILI), acute respiratory infection (ARI) or both to a national focal point. From the national level, both numerator and denominator data are then reported to the European Surveillance System (TESSy) database. Additional semi-quantitative indicators of intensity, geographic spread and trend of influenza activity at the national level are also reported.

Virological surveillance

Weekly analysis – virology

In week 09/2010, 22 countries and the UK (England, Northern Ireland and Scotland) reported virological data. Sentinel physicians collected 425 specimens, 20 (4.7%) of which were positive for influenza virus (Tables 1 and 2). In addition, 129 non-sentinel source specimens (i.e. specimens collected for diagnostic purpose in hospitals) were reported positive for influenza virus. Of the 149 influenza viruses detected from sentinel and non-sentinel sources during week 9/2010, 19 (13%) were type B viruses. Of the 16 151 type A influenza viruses detected by sentinel practices and on which sub-typing was performed since week 40/2009, 16 094 (99%) were identified as the 2009 pandemic influenza A(H1N1) virus. Table 2 shows the distribution of both sentinel and non-sentinel specimens by type and sub-type. Figures 1–3 show the trends of virological detections over time. The proportion of positive sentinel samples has decreased since week 46/2009 (Figure 3).

More details on circulating viruses can be found in the [report](#) prepared by the Community Network of Reference Laboratories (CNRL) coordination team.

From week 40/2009 to week 09/2010, 3027 antigenic and/or genetic characterisations were reported, of these 2974 (98%) were A/California/7/2009(H1N1)-like, six (<1%) were A/Brisbane/10/2007(H3N2)-like, 39 (1%) were A/Perth/16/2009(H3N2)-like, seven (<1%) were B/Brisbane/60/2008-like (B/Victoria/2/87 lineage) and one (<1%) was A/Brisbane/59/2007 (H1N1)-like. The antigenic characterisations of the influenza viruses from week 40/2009 to week 9/2010 reflect the same pattern (Table 3).

All pandemic viruses tested were resistant to M2 inhibitors. Of the 1453 viruses tested from nine countries, 37 (2.5%) were resistant to oseltamivir, and of the 1447 viruses tested, none were resistant to zanamivir (Table 4).

Since the peak in week 01/2010, the total number of respiratory syncytial virus (RSV) detections in 11 countries has been decreasing (Figure 4).

Table 2: Weekly and cumulative influenza virus detections by type, subtype and surveillance system, weeks 40/2009–09/2010

Virus type/subtype	Current Week		Season	
	Sentinel	Non-sentinel	Sentinel	Non-sentinel
Influenza A	15	115	16808	89761
A (pandemic H1N1)	13	94	16094	78191
A (subtyping not performed)	2	21	657	11431
A (not subtypable)	0	0	14	47
A (H3)	0	0	8	43
A (H1)	0	0	35	49
Influenza B	5	14	74	138
Total Influenza	20	129	16882	89899

Note: A(pandemic H1N1), A(H3) and A(H1) includes both N-subtyped and not N-subtyped viruses

Figure 1: Number of sentinel specimens positive for influenza, by type, subtype and by week of report, weeks 40/2009–09/2010




Figure 2: Number of non-sentinel specimens positive for influenza by type, subtype and week of report, weeks 40/2009–09/2010




Figure 3: Proportion of sentinel samples positive for influenza, weeks 40/2009–09/2010





Table 3: Results of antigenic characterisations of sentinel and non-sentinel influenza virus isolates since week 40/2009

Strain name	Number of strains
A/California/7/2009 (H1N1)-like	1915
A/Brisbane/10/2007 (H3N2)-like	5
A/Perth/16/2009 (H3N2)-like	22
B/Brisbane/60/2008-like (B/Victoria/2/87 lineage)	4

Table 4: Antiviral resistance by influenza virus type and subtype, weeks 40/2009–09/2010

Virus type and subtype	Resistance to neuraminidase inhibitors				Resistance to M2 inhibitors	
	Oseltamivir		Zanamivir		Isolates tested	Resistant n (%)
	Isolates tested	Resistant n (%)	Isolates tested	Resistant n (%)		
A(H3N2)	0	0	0	0	0	0
A(H1N1)	0	0	0	0	0	0
A(H1N1)v	1453	37 (2.5%)	1447	0 (0%)	205	205 (100%)
B	0	0	0	0	0	0

Figure 4: Respiratory syncytial virus (RSV) detections, sentinel and non-sentinel, weeks 40/2009–09/2010



Description of the system

According to the nationally defined sampling strategy, sentinel physicians take nasal or pharyngeal swabs from patients with influenza-like illness (ILI), acute respiratory infection (ARI) or both and send the specimens to influenza-specific reference laboratories for virus detection, (sub-)typing, antigenic or genetic characterisation and antiviral susceptibility testing.

For details on the current virus strains recommended by WHO for vaccine preparation [click here](#).

Aggregate numbers of pandemic (H1N1) 2009 deaths

Weekly analysis — deaths

During week 9/2010, six countries reported 6 deaths. Since the beginning of the pandemic, 1819 deaths have been notified to ECDC through TESSy (Table 5).

Table 5: Aggregate numbers of pandemic (H1N1) 2009 deaths, week 09/2010

country	Deaths reported in week	Cumulative deaths since start of season	Last reported week
Austria		0	2009-w36
Belgium		0	2009-w29
Bulgaria		40	2009-w53
Cyprus		0	2009-w29
Czech Republic	0	97	2010-w09
Denmark		0	2009-w36
Estonia	1	19	2010-w09
Finland		0	2009-w36
France	1	308	2010-w09
Germany	0	243	2010-w09
Greece	1	138	2010-w09
Hungary	1	130	2010-w09
Iceland		2	2009-w52
Ireland	1	23	2010-w09
Italy		1	2009-w52
Latvia	0	34	2010-w09
Lithuania	0	23	2010-w09
Luxembourg		3	2009-w52
Malta		5	2010-w08
Netherlands	0	58	2010-w09
Norway		29	2010-w08
Poland		148	2009-w53
Portugal		0	2009-w36
Romania	0	122	2010-w09
Slovakia	1	53	2010-w09
Slovenia		19	2010-w07
Spain		4	2009-w29
Sweden	0	24	2010-w09
United Kingdom	0	296	2010-w09
Total	6	1819	

Description of the system

Aggregate numbers of both probable and laboratory-confirmed cases of pandemic influenza and deaths due to pandemic influenza are reported by countries still collecting this data. As countries are retrospectively updating their weekly numbers of deaths and the system calculates the cumulative values based on the current status, weekly numbers of deaths published in previous WISO editions may not always add up to the cumulative totals.

Hospital surveillance – severe acute respiratory infection (SARI)

Weekly analysis – SARI

During week 09/2010, five SARI cases were reported and none of them had symptom onset during the same week. The number of SARI cases by week of onset has been in decline since the peak in week 46/2009 (Figure 5). Since the beginning of SARI surveillance, 11 countries have reported 11 295 cases, including 554 fatalities (Table 6).

More than 99% of the influenza viruses isolated from SARI cases since the start of the season were the 2009 pandemic influenza virus (Table 7).

Table 6: Cumulative number of SARI cases, weeks 40/2009 - week 09/2010

Country	Number of cases	Incidence of SARI cases per 100,000 population	Number of fatal cases reported	Incidence of fatal cases per 100,000 population	Estimated population covered
Austria	2821		38		
Belgium	1865	17.48			10668666
Cyprus	25		8		
Finland	1421	26.68	55	1.03	5326314
France	1339		292		
United Kingdom	1567	3.97	64	0.16	39503332
Ireland	903		17		
Malta	181	43.76	1	0.24	413609
Netherlands	645	3.9	27	0.16	16521505
Romania	197	1.55	12	0.09	12684180
Slovakia	331		40		
Total	11295		554		85117606

Figure 5: Number of SARI cases by week of onset, week 09/2010




Table 7: Number of SARI cases by influenza type and subtype, week 09/2010

Virus type/subtype	Number of cases during current week	Cumulative number of cases since the start of the season
Influenza A	1	8983
A (pandemic H1N1)	1	8953
A(subtyping not performed)		23
A(H3)		
A(H1)		7
A(H5)		
Influenza B		
Unknown	4	2312
Total	5	11295

Description of the system

A number of Member States carry out hospital-based surveillance of severe acute respiratory infection (SARI) exhaustively or at selected sentinel sites. SARI surveillance serves to monitor the trends in the severity of influenza and potential risk factors for severe disease to help guide preventive measures and health care resource allocation.

Qualitative reporting

Qualitative monitoring will be an acceptable replacement for the quantitative monitoring when reliable numbers are no longer available for reporting due to overburdened surveillance systems. The qualitative components will give some indication of influenza intensity, geographic spread, trend and impact.

Erratum

On page six of the Weekly Influenza Surveillance Overview for week 6, it was reported that the total number of pandemic viruses tested for resistance to oseltamivir was 1451. The correct number was 1453.

The report text was written by an editorial team at the [European Centre for Disease Prevention and Control](#) (ECDC): Flaviu Plata, Phillip Zucs, Bruno Ciancio, Rene Snacken and Eeva Broberg. The bulletin text was reviewed by the Community Network of Reference Laboratories for Human Influenza in Europe (CNRL) coordination team: Adam Meijer, Rod Daniels, Alan Hay and Maria Zambon. On behalf of the EISN members the bulletin text was reviewed by Joan O'Donnell (Health Protection Surveillance Centre, Ireland) and Katarina Prosenc (National Institute of Public Health, Slovenia).

Maps and commentary used in this Weekly Influenza Surveillance Overview (WISO) do not imply any opinions whatsoever of ECDC or its partners on the legal status of the countries and territories shown or concerning their borders.

All data published in the WISO are up-to-date on the day of publication. Past this date, however, published data should not be used for longitudinal comparisons as countries tend to retrospectively update their numbers in the database.

© European Centre for Disease Prevention and Control, Stockholm, 2009